

ALDERFER & TRAVIS CARDIOLOGY, PC
670 Lawn Ave., Suite 3A Sellersville, PA 18960
Tel (215)257-9500 FAX (215) 257-9500

STREPTOKINASE - INJECTION

USES: This medication is an enzyme which works to break up and dissolve blood clots which can block arteries. It is used in the treatment of heart attack, pulmonary embolism or blood clots. It may also be used to clear blood clots from catheters and other devices.

HOW TO USE THIS MEDICATION: This medication is given by injection by a health care professional.

It is important this medication be used as prescribed. It is most effective when administered as soon as possible (within 6 hours after symptoms (e.g., chest pain) appear.

SIDE EFFECTS: Nausea, dizziness, low blood pressure or mild fever may occur. It can also cause nerve damage. If any of these effects continue or become bothersome, inform your doctor.

Notify your doctor if you experience easy bruising or bleeding, a skin rash, breathing trouble, headache, flushing, itching, a rapid or abnormal heartbeat or chest pain while using this medication.

Immediately report any signs of bleeding to your doctor.

PRECAUTIONS: Tell your doctor if you have any pre-existing bleeding disorders, high blood pressure, endocarditis, recent biopsy or surgery, recent injury or any allergies.

Use extra caution to avoid injury and trauma (e.g., carefully brush teeth) while using this medication due to the increased risk of bleeding.

This medication should be used only if clearly needed during pregnancy. Discuss the risks and benefits with your doctor. It is not known if this medication appears in breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS: Tell your doctor of any over-the-counter or prescription medication you may take including any blood thinners, NSAID or aspirin.

Drugs that can reverse effects of streptokinase include aminocaproic acid, aprotinin and tranexamic acid.

NOTES: Laboratory tests will be done frequently while using this medication to monitor its effects and to prevent side effects.

MISSED DOSE: Continued or repeated doses are usually not

necessary for this medication.

STORAGE: Store in a cool area as directed. Discard any unused medication after the expiration date on the label. Reconstituted solution may be refrigerated between 36 to 46 degrees F (2 to 8 degrees C) if used within 8 hours. Discard unused portion.